

Product-Data-Sheet for PODO/SVTERT152

Version: January 2020

For questions please contact office@evercyte.com

Evercyte Ord. No.:	CLHT-033-0152
Designation:	PODO/SVTERT152
Biosafety Level:	1
Shipped:	Frozen on dry ice
Medium:	<p>PodUp3 (Evercyte, Cat# MHT-033-3): MCDB131 Basal Medium (Pan Biotech, Cat# P04-80057) 1,6 mM GlutaMAX-I (Gibco, Cat# 35050-038) 9,6 µg/mL Bovine Brain Extract (Lonza, Cat# CC-4098) 8 ng/ml hEGF (Sigma Aldrich, Cat# E9644) 20 ng/ml Hydrocortisone (Sigma Aldrich, Cat# H0396) 20% fetal bovine serum (Sigma Aldrich, Cat# F7524) 100 µg/ml G418 (InvivoGen, Cat# ant-gn-5)</p>
Growth:	Adherent
Organism:	Homo sapiens (human)
Morphology:	Typical podocyte-like, cytoplasmic extensions with arborized appearance
Source:	Human urine sediment
Cell Type:	Podocytes
Antigen Expression:	Positive for nephrin, WT-1, synaptopodin
Ethical statement:	Approved by Institutional Review Board (IRB) in accordance with the Declaration of Helsinki.

Comments:	<p>PODO/SVTERT152 was developed from human urine-derived podocytes by transfection with a plasmid encoding SV40 early region followed by transduction with a retroviral expression vector (pLXSN) containing the hTERT gene.</p> <p>Sub-confluent cultures show flat cytoplasmatic protrusions and occasional elongated processes. Within the culture rare interspersed multinuclear cells with an arborized appearance are present. Confluent cultures acquire a more cobblestone morphology.</p> <p>The cells readily recover from cryopreservation and can be continuously cultured for a minimum of 50 population doublings after thawing. No changes in growth characteristics have been observed after thawing.</p>
Propagation:	<p>Cells are grown in PodoUp3 medium (see above) at 37°C in a humidified atmosphere with 5 % CO₂.</p>
Subculturing:	<p>New culture flasks have to be pre-coated with human collagen I. Therefore, the culture flasks are pre-treated with collagen I solution (Sigma Aldrich, Cat# C2249-20ML, diluted to 50 µg/mL in PBS; 70 µl/cm²) at 37°C for at least 30 min. Before introducing cells, remove excess of collagen I solution and rinse flask once with PBS (160 µl/cm²).</p> <p>For detachment of the cells remove and discard the culture medium and wash the cells twice with PBS. Remove PBS completely. Then, add 0.05% Trypsin-EDTA (1x) solution (room-temperature; 20 µl/cm²; Gibco, Cat# 25300054), make sure that all cells have been in contact with this solution and incubate the culture flask at 37°C for approximately 3 – 4 min. Observe cell detachment under an inverted microscope. As soon as all cells are detached (if necessary agitate the cells by gently hitting the flask), add growth medium (about 160 µl/cm²) and aspirate cells by pipetting. Add appropriate aliquots of the cell suspension to collagen I pre-coated culture vessels supplemented with growth medium (final volume of 240 µl/cm²).</p> <p>Cells should be split every 3-4 days (<u>after having reached not more than 80 % confluence</u>) with a split-ratio of 1:6 to 1:8.</p>
Preservation:	<p>Freezing medium: PodoUp3 (Evercyte, Cat# MHT-033) with 10 % DMSO Storage temperature: liquid nitrogen</p>
Freezing and thawing procedure:	<p>Freezing of cells: Detach the cells from the culture vessel by using Trypsin-EDTA solution as described above, resuspend the detached cells in growth medium and centrifuge at 170 g for 5 min. Then, discard the supernatant, resuspend the resulting cell pellet in the remaining droplet and add freezing medium (tempered to 4°C) to reach a cell density of about 8 x 10⁵ cells/ml (for thawing in a 25 cm² culture flask). Add 1 ml of this cell suspension to each pre-cooled cryovial and immediately transfer the cells to -80°C.</p>

	<p>After 24 hours transfer the vials to the liquid nitrogen tank.</p> <p>Thawing of cells: Pre-coat a 25 cm² culture flask with collagen I (see subculturing). Then, add 6 ml of growth medium to the prepared culture flask and transfer it to the incubator for at least 20 min to allow the medium to reach its normal pH. Take a vial of frozen cells, rinse outside with Ethanol and pre-warm in hand until one last piece of frozen cells is seen. Then, immediately transfer the content of the vial to a 15 ml centrifugation tube pre-filled with 9 ml of medium pre-cooled to 4°C and centrifuge for 5 min at 170 g. Then, discard the supernatant and resuspend the cell pellet in the remaining droplet. Add 1 ml of pre-warmed medium to the cells, transfer the cells to the prepared culture flask and incubate at 37°C in a suitable incubator. Perform a medium change 24 hours after thawing. If the cells are near confluent at this point, they should be passaged (see subculturing).</p>
Doubling Time:	About 24-32 hours
Virus Testing:	Cells have been tested negative for HAV and Parvo B19 with Roche DPX-PCR (cobas [®] TaqScreen DPX-Test), for HBV, HCV, HIV nucleic acids with Roche-Multiplex-PCR (cobas [®] TaqScreen MPX Test, v2.0).
Other Analytical Data:	<p>Cells are negative for Mycoplasma contaminations as tested using MycoAlert[™] Mycoplasma Detection Kit from Lonza.</p> <p>Cells are negative for bacterial and fungal contaminations as tested according Ph. Eur. 2.6.1. / USP <71>.</p> <p>STR profile has been analysed and is as expected.</p>

Please Note:

The classification of biosafety level is based on the directive 2000/54/EG of the European Parliament and of the Council on the protection of workers from risks related to exposure to biological agents at work. While Evercyte undertakes all reasonable measures to test for absence of a selected panel of known human pathogenic viruses, there is currently no test procedure available that guarantees for complete absence of infectious pathogens. The use of state-of-the art infectious virus assays or viral antigen assays may leave open the possible existence of a latent viral genome, even if a negative test result is obtained. Therefore, we recommend that all human cell lines should be handled with caution such as an organism of ACDP Hazard Group 2.

Notices and disclaimers: Evercyte products are intended for laboratory research purposes only. They are not intended for use in humans.

While Evercyte uses reasonable efforts to include accurate and up-to-date information on this site, no warranties or representations are made to its accuracy. Citations from scientific literature and patents are provided for informational purposes only and no warranties are made that such information has been confirmed to be accurate.